

On the cosmological distance and redshift between any two objects

J. Liske[★]

School of Physics, University of New South Wales, Sydney 2052, Australia

Accepted Received

ABSTRACT

We discuss the problem of how to calculate the distance between two cosmological objects given their redshifts and angular separation on the sky. Although of a fundamental nature, this problem and its solution seem to lack a detailed description in the literature. We present a new variant of its solution and quantitatively assess the most commonly used approximation.

Key words: cosmology: miscellaneous

1 INTRODUCTION

In cosmology and extragalactic astronomy one frequently needs to calculate the distance between two objects given their redshifts and their angular separation on the sky. As larger and larger cosmological volumes are probed by wide field redshift surveys such as 2dF and the Sloan Digital Sky Survey, or by QSO absorption line studies, the effects of non-Euclidean geometry become increasingly important. In the past, authors have frequently relied on approximations when calculating the distance between two cosmological objects, presumably for calculative ease. Such approximations are valid only for small distances and are particularly useful when examining or highlighting the feasibility of geometrical means to measure cosmological parameters as was done e.g. by Alcock & Paczyński (1979), Phillipps (1994) and Popowski et al. (1998). However, many practical applications are not limited by computing time and since an exact solution to the problem exists the approximations seem unnecessary. These applications include the construction of the real-space two-point correlation function of various objects such as galaxies (e.g. Yoshii, Peterson, & Takahara 1993) and QSO absorption systems (Williger et al. 1996; Dinshaw & Impey 1996), as well as studies of the effect of local sources of ionizing radiation on their surrounding intergalactic medium (e.g. Fernández-Soto et al. 1995). The latter application actually requires knowledge of not only the distance between two objects but also of the redshift experienced by a photon travelling from one object to another.

Given the fundamental nature of this problem we feel a detailed, explicit treatment is called for. In this short article we present a new variant of the solution to the distance problem (Sections 2 and 3), discuss its relation to existing

approaches (Section 4) and finally investigate the validity of the most frequently used approximation (Section 5).

For clarity and brevity we have limited ourselves in this paper to homogeneous Friedmann (zero-pressure) cosmologies with no cosmological constant ($\Lambda = 0$). The inclusion of Λ renders some of the explicit expressions non-analytical and thus (in the context of this paper) unnecessarily complicates matters.

2 THE DISTANCE BETWEEN ANY TWO OBJECTS

We begin by writing the familiar Robertson-Walker line element as:

$$ds^2 = -c^2 dt^2 + a^2(t) [d\chi^2 + \Sigma^2(\chi)(d\theta^2 + \sin^2 \theta d\phi^2)], \quad (1)$$

where

$$\Sigma(\chi) = \begin{cases} \sin \chi & k = +1 \\ \chi & k = 0 \\ \sinh \chi & k = -1. \end{cases} \quad (2)$$

Putting the Earth at the origin of the coordinate system one can use this metric and the Friedmann equations to calculate the distance from Earth (at $\chi = 0$) to an object at redshift z , corresponding to a comoving coordinate χ ,

$$\begin{aligned} r &= a_0 \Sigma[\chi(z)] \\ &= \frac{c}{H_0 q_0^2} \frac{1}{1+z} \\ &\quad \times \left[q_0 z + (q_0 - 1) \left(\sqrt{1 + 2q_0 z} - 1 \right) \right] \end{aligned} \quad (3)$$

(e.g. Misner, Thorne, & Wheeler 1973), where a_0 , H_0 and q_0 are the scale factor, Hubble and deceleration parameters at the present epoch (subscript 0).

[★] E-mail: jol@phys.unsw.edu.au

Now consider an object 1 (the ‘receiver’) observed on Earth today at z_1 and an object 2 (the ‘emitter’) at z_2 separated by an angle α on the sky (cf. Fig. 1). Object 2 emits a photon towards object 1 which is received by object 1 at the same time as object 1 emits the photon we receive from it today (i.e. at the epoch corresponding to z_1). What is the distance, r'_2 , between these two objects at the time of the photon reception and what is the redshift, z'_2 , of the photon as observed by object 1?

An observer at object 1 would write equation (3) as

$$\begin{aligned} r'_2 &= a_1 \Sigma(\chi'_2) \\ &= \frac{c}{H_1 q_1^2} \frac{1}{1+z'_2} \\ &\quad \times \left[q_1 z'_2 + (q_1 - 1) \left(\sqrt{1 + 2q_1 z'_2} - 1 \right) \right], \end{aligned} \quad (4)$$

where χ'_2 is the comoving coordinate distance between objects 1 and 2 (cf. Fig. 1) and

$$\begin{aligned} a_1 &= \frac{a_0}{1+z_1}, \\ H_1 &= H_0(1+z_1)\sqrt{1+2q_0 z_1}, \\ q_1 &= q_0 \frac{1+z_1}{1+2q_0 z_1} \end{aligned} \quad (5)$$

are the scale factor, Hubble and deceleration parameters at the time object 1 emitted the photons we receive today. Thus we see that the problems of calculating r'_2 and z'_2 are equivalent since knowledge of one provides knowledge of the other via equation (4). Here we choose to find z'_2 .

Note that r'_2 does *not* describe in general the shortest distance between objects 1 and 2 along a $t = \text{const}$ hypersurface of spacetime. Nevertheless, in many applications r'_2 is the quantity of interest. E.g. when considering the radiative effect of a QSO on a nearby object one needs the luminosity distance between the two which is given by $r'_{L,2} = r'_2(1+z'_2)$. In any case, it is most practical to solve the problem for z'_2 and then calculate the distance required for a given application from z'_2 .

As we are dealing with three points (Earth and objects 1 and 2) in a 3-dimensional space (described by the spatial part of the metric, equation 1) it seems intuitive that it must be possible to reduce the problem to two dimensions as one can always find a 2d hypersurface that contains all three points. Since we are interested in measuring 3d distances the hypersurface should be chosen such that the distance between any two points of the surface (as measured along the surface) is identical to the 3d distance between the same two points. We call such surfaces totally geodesic. It is clear that a totally geodesic hypersurface containing a given set of three points can only be constructed from the geodesics connecting the three points. Since these geodesics are unique there can be only one such surface.

In equation (1) we introduced a polar coordinate system (χ, θ, ϕ) . Clearly, the hypersurface described by $\phi = \text{const}$ is totally geodesic. (Note that the surfaces $\chi = \text{const}$ and $\theta = \text{const}$ are not.) Since the curvature of the 3d space under consideration is constant, one can generate *all* totally geodesic hypersurfaces from any given one by mere translations and rotations. Therefore, for a given set of three points there must exist a coordinate system $(\tilde{\chi}, \tilde{\theta}, \tilde{\phi})$ such that $\tilde{\phi} = \text{const}$ describes the unique totally geodesic hypersurface containing these three points. Since this new coordi-




Figure 1. In the case $k = +1$ objects 1 and 2 and Earth form a triangle with geodesic sides (thick lines) on the surface of a 2d sphere of radius a . The Earth is at the origin of the coordinate system (‘north pole’). From the centre of the sphere the angle between Earth and object 1 is χ_1 . A photon emitted by object 1 towards Earth travels along the geodesic connecting the two (thick line). The length of this path (= distance between object 1 and Earth) is $a\chi_1$. $a\Sigma(\chi_1) = a \sin \chi_1$ is the distance from object 1 to the central axis of the sphere. Essentially this is the angular diameter distance or luminosity distance (modulo factors of $1+z$) from Earth to object 1. χ'_2 is the unknown angle between objects 1 and 2 at the centre of the sphere and $a\chi'_2$ is the unknown length of the connecting geodesic.

nate system can be constructed from the old one by translation and rotation, the form of the metric in this new system is identical to equation (1). Restraining this metric to the $\tilde{\phi} = \text{const}$ hypersurface we have

$$dl^2 = a^2 [d\tilde{\chi}^2 + \Sigma^2(\tilde{\chi})d\tilde{\theta}^2]. \quad (6)$$

Thus we can see that the triangle Earth–object 1–object 2 lies either on a 2d sphere, a plane or a 2d hyperboloid ($k = +1, 0, -1$) embedded in 3d Euclidean space. This triangle has geodesic sides χ_1, χ_2 , and χ'_2 (connecting the two objects) and the angle α at Earth. The case $k = +1$ is shown in Fig. 1.

The objective is now to express the unknown side χ'_2 in terms of the known sides χ_1, χ_2 and the angle α . We first note that all formulae of Euclidean trigonometry have corresponding formulae in spherical and hyperbolic trigonometry. These can be expressed simultaneously for all three curvatures using Σ . In particular, we can generalize the half-angle formulae in this way and use them to show that

$$\begin{aligned} \Sigma^2 \left(\frac{\chi'_2}{2} \right) &= \Sigma^2 \left(\frac{\chi_2 + \chi_1}{2} \right) \sin^2 \frac{\alpha}{2} \\ &\quad + \Sigma^2 \left(\frac{\chi_2 - \chi_1}{2} \right) \cos^2 \frac{\alpha}{2}. \end{aligned} \quad (7)$$

This is a more compact and symmetrical version of the generalized cosine rule (see Section 4).

Using the same methods that were employed in the derivation of equation (3) we can relate the right-hand side of equation (7) to z_1 and z_2 :

$$a_0 \Sigma \left(\frac{\chi_2 \pm \chi_1}{2} \right) = \frac{c}{H_0 q_0} \frac{1}{\sqrt{(1+z_1)(1+z_2)}} \frac{P_{\pm}}{2}, \quad (8)$$

where

$$P_+ = \frac{1}{q_0} \left[(q_0 - 1) \left(\sqrt{1+2q_0 z_1} + \sqrt{1+2q_0 z_2} - 1 \right) + \sqrt{(1+2q_0 z_1)(1+2q_0 z_2)} - q_0 \right] \quad (9)$$

and

$$P_- = \left(\sqrt{1+2q_0 z_2} - \sqrt{1+2q_0 z_1} \right). \quad (10)$$

Furthermore, setting $\chi_1 = z_1 = 0$ it follows from equation (8) that

$$a_0 \Sigma \left(\frac{\chi_2}{2} \right) = \frac{c}{H_0 q_0} \frac{1}{\sqrt{1+z_2}} \frac{1}{2} \left(\sqrt{1+2q_0 z_2} - 1 \right) \quad (11)$$

and for an observer at object 1

$$a_1 \Sigma \left(\frac{\chi'_2}{2} \right) = \frac{c}{H_1 q_1} \frac{1}{\sqrt{1+z'_2}} \frac{1}{2} \left(\sqrt{1+2q_1 z'_2} - 1 \right), \quad (12)$$

thus relating the left-hand side of equation (7) to z'_2 . Solving

$$P = \frac{1}{2} \frac{1}{\sqrt{1+z'_2}} \left(\sqrt{1+2q_1 z'_2} - 1 \right) \quad (13)$$

for z'_2 yields (positive solution)

$$z'_2 = \frac{2P^2}{(q_1 - 2P^2)^2} \left(1 + q_1 - 2P^2 + \sqrt{\frac{q_1^2}{P^2} + 1 - 2q_1} \right), \quad (14)$$

where

$$\begin{aligned} P^2 &= \left(\frac{a_1 H_1 q_1}{c} \right)^2 \Sigma^2 \left(\frac{\chi'_2}{2} \right) \\ &= \frac{1}{4} \frac{1+z_1}{1+z_2} \frac{1}{1+2q_0 z_1} \left(P_+^2 \sin^2 \frac{\alpha}{2} + P_-^2 \cos^2 \frac{\alpha}{2} \right) \end{aligned} \quad (15)$$

Note that z'_2 does not depend on H_0 .

2.1 $\Lambda \neq 0$

For completeness we now briefly consider the case $\Lambda \neq 0$. Equation (7) is of course still valid. However, in relating this equation to the observables z_1 , z_2 and z'_2 we have made use of the fundamental relationship between an object's comoving radial coordinate and its redshift,

$$\chi(z) = \frac{c}{a_0 H_0} \int_1^{1+z} \left[\Omega_0 x^3 + (1 - \Omega_0) x^2 \right]^{-\frac{1}{2}} dx, \quad (16)$$

which in turn is derived from the Friedmann equations. For $\Lambda \neq 0$ these take on a different structure and instead of the above relationship we have

$$\chi(z) = \frac{c}{a_0 H_0} \int_1^{1+z} \left[\Omega_0 x^3 + (1 - \Omega_0 - \lambda_0) x^2 + \lambda_0 \right]^{-\frac{1}{2}} dx, \quad (17)$$

where $\lambda_0 = \Lambda c^2 / 3H_0^2$. Unfortunately this integral is non-analytical but Kayser, Helbig, & Schramm (1997) developed an efficient method (which also accommodates inhomogeneity) to compute $\Sigma(\chi)$ numerically.

Note that other density contributions with unusual equations of state can be dealt with in the same way. All that needs to be done is to establish $\chi(z)$ or, equivalently,




Figure 2. z'_2 as a function of α (angle on the sky between receiver and emitter), $z_1 = z_2$. Thick lines are for $q_0 = 0.5$, thin lines for $q_0 = 0.15$.




Figure 3. z'_2 as a function of $z_1 = z_2$. Thick lines are for $q_0 = 0.5$, thin lines for $q_0 = 0.15$.

$\Sigma[\chi(z)]$. In principle, one could then proceed to use equation (7) to find z'_2 and r'_2 . However, when $\chi(z)$ is not available analytically it is probably more practical to use the generalized cosine rule (see Section 4, equation 23) instead of equation (7). In any case, one is still left with the problem of inverting $\Sigma[\chi(z)]$ in order to find z'_2 . However, since we can compute

$$\frac{d\Sigma}{dz} = \sqrt{1 - k\Sigma^2(z)} \frac{d\chi}{dz} \quad (18)$$

it should be possible to employ an efficient root finding algorithm for this task.

3 RESULTS

As cosmologists are often used to thinking in terms of redshift rather than distance, we show the result of the above calculations in Figs. 2–6 in terms of z'_2 , the redshift of object 2 as seen by object 1.

Figs. 2 and 3 explore the special case $z_1 = z_2$. In Fig. 2 we show $z'_2(\alpha)$ with $z_1 = z_2$ fixed at various values. The thick lines show the case of a flat universe ($\Omega_0 = 2q_0 = 1$) and the




Figure 4. z'_2 as a function of z_1 (redshift of receiver), $\alpha = 0$. Thick lines are for $q_0 = 0.5$, thin lines for $q_0 = 0.15$. Since $\alpha = 0$, z'_2 is independent of cosmology for $z_1 < z_2$.

thin lines show the case of an open universe with $\Omega_0 = 0.3$. Note that for large values of $z_1 = z_2$ there is some α_∞ such that $z'_2 \rightarrow \infty$ for $\alpha \rightarrow \alpha_\infty$. This is the particle horizon of object 1 at the epoch corresponding to z_1 . At that time, light emitted from objects separated from object 1 by angles $> \alpha_\infty$ has not had time to reach object 1 since the Big Bang (ignoring inflation). In Fig. 3 we fix α at various angles and show z'_2 as a function of $z_1 = z_2$. The case $\alpha = 180^\circ$ is often incorrectly used in undergraduate physics textbooks (e.g. Halliday, Resnick, & Walker 1993, p. 1128–1129) as an example of how to add velocities in Special Relativity, a method which will give a wrong result for z'_2 .

In Figs. 4 and 5 we consider the special case $\alpha = 0$ and plot z'_2 as a function of z_1 and z_2 respectively. Thick lines again represent a flat universe, thin lines an open universe. Whenever $z_1 < z_2$ (and $\alpha = 0$), z'_2 is of course given by

$$1 + z'_2 = \frac{1 + z_2}{1 + z_1} \quad (19)$$

which is the only case where z'_2 is independent of the cosmological model, since in this case the time of emission of the photon received by object 1 is the same as the time corresponding to z_2 . Note that although $\chi'_2(z_1, z_2, \alpha) = \chi'_2(z_2, z_1, \alpha)$ (cf. equation 23), the same does *not* hold for z'_2 (cf. equations 14 and 15). Fig. 5 also provides the solution to an interesting thought experiment. What is the redshift, z_{refl} , of a photon emitted by ourselves ($z_2 = 0$) which was reflected back to us by a comoving mirror at z_1 ? The answer is

$$1 + z_{\text{refl}} = (1 + z_1)[1 + z'_2(z_1, z_2 = 0)]. \quad (20)$$

Finally we plot in Fig. 6 lines of constant z'_2 as a function of z_2 and α for a receiver at $z_1 = 3$. In this plot the flat and open cosmologies are represented by the solid and dashed lines respectively.

4 RELATION TO OTHER SOLUTIONS

We are aware of two original solutions to the problem in the literature, Peacock (1999), p. 71, and Osmer (1981). Neither of them consider z'_2 .




Figure 5. z'_2 as a function of z_2 (redshift of emitter), $\alpha = 0$. Thick lines are for $q_0 = 0.5$, thin lines for $q_0 = 0.15$. Since $\alpha = 0$, z'_2 is independent of cosmology for $z_1 < z_2$.




Figure 6. z'_2 as a function of z_2 and α (emitter position). The receiver is at $z_1 = 3$ and $\alpha = 0$ (marked by a diamond). The contours are lines of constant z'_2 . Solid lines are for $q_0 = 0.5$, dashed lines for $q_0 = 0.15$.

Instead of equation (7), Peacock considers the generalized cosine rule which directly gives $\Sigma(\chi'_2)$. Introducing the cosine equivalent of $\Sigma(\chi)$,

$$\Xi(\chi) = \sqrt{1 - k\Sigma^2(\chi)}, \quad (21)$$

we have

$$\Xi(\chi'_2) = \Xi(\chi_1) \Xi(\chi_2) + k\Sigma(\chi_1)\Sigma(\chi_2) \cos \alpha \quad (22)$$

which can be written as

$$\begin{aligned} \Sigma^2(\chi'_2) &= \Sigma^2(\chi_1) \Xi^2(\chi_2) + \Sigma^2(\chi_2) \Xi^2(\chi_1) \\ &\quad + k\Sigma^2(\chi_1)\Sigma^2(\chi_2) \sin^2 \alpha \\ &\quad - 2\Sigma(\chi_1)\Sigma(\chi_2) \Xi(\chi_1) \Xi(\chi_2) \cos \alpha. \end{aligned} \quad (23)$$

When only r'_2 is needed the use of this equation seems more practical than our solution presented in Section 2. However, when using the generalized cosine rule the analogues of equations (14) and (15) are more complicated so that in cases where z'_2 is (also) of interest, e.g. when calculating the luminosity distance $r'_{12} = r'_2(1 + z'_2)$, the new solution is to be preferred.




Figure 7. The contours are lines of constant $\frac{a_0 \Delta \chi'_2}{a_0 \chi'_2}$, the fractional error made when using equation (25) for $z_1 = 3$ (marked by a diamond) and $q_0 = 0.15$. The contour levels are indicated.

Osmer's solution may be considered the most rigorous as it is based on a general result of differential geometry in maximally symmetric spaces. Weinberg (1972), p. 413, showed how to transform to a coordinate system which has been 'quasitranslated'. Osmer uses this equation to transform from a coordinate system in which objects 1 and 2 have particularly simple coordinates (see discussion in Section 2) to one where the origin has been translated from Earth to object 1. The result is then given by

$$\Sigma^2(\chi'_2) = \Sigma^2(\chi_2) \sin^2 \alpha + [\Sigma(\chi_2) \Xi(\chi_1) \cos \alpha - \Sigma(\chi_1) \Xi(\chi_2)]^2. \quad (24)$$

This equation of course reduces to equation (23) and thus the comments made there apply equally to Osmer's solution.

5 VALIDITY OF APPROXIMATION

Probably the most common approximation for the comoving distance is (e.g. Yoshii, Peterson, & Takahara 1993; Phillipps 1994)

$$a_0^2 \chi'^2_2 \approx a_0^2 \chi'^2_{A2} = a_0^2 \Sigma^2[\chi(\bar{z})] \alpha^2 + \left[a_0 \frac{d\chi}{dz}(\bar{z}) \right]^2 \Delta z^2, \quad (25)$$

where $\Delta z = z_1 - z_2$ and $\bar{z} = (z_1 + z_2)/2$ and

$$a_0 \frac{d\chi}{dz}(z) = \frac{c}{H(z)}. \quad (26)$$

In Fig. 7 we plot the fractional error made when using equation (25),

$$\frac{a_0 \Delta \chi'_2}{a_0 \chi'_2} = \frac{a_0 \chi'_{A2} - a_0 \chi'_2}{a_0 \chi'_2}, \quad (27)$$

against z_2 and α for object 1 at $z_1 = 3$ and for $q_0 = 0.15$. We see that the approximation gives both too large and too small distances (solid and dashed contours) depending on the position in the z_2 - α plane. Note the 'ridges' along which the approximation incidentally gives the correct distance.

The special case $\Delta z = 0$ deserves some further attention as it corresponds to the well-known angular diameter problem: what is the length, L , of a rod at z which subtends

an angle α as seen from Earth? Commonly, the answer is given as

$$L_A = \alpha r_A = \frac{\alpha a_0 \Sigma(\chi)}{1+z} \quad (28)$$

where r_A is the angular diameter distance. However, strictly speaking L_A is the length of the line $\chi = \text{const}$ connecting the two ends of the rod at the epoch z , which is *not* the shortest distance between the ends. The length of the geodesic connecting the two ends is given by $a\chi'_2$ which we can derive from equation (7). Setting $\chi_1 = \chi_2 = \chi$ and $z_1 = z_2 = z$ we arrive at

$$\begin{aligned} L &= a\chi'_2 \\ &= \frac{a_0}{1+z} 2 \Sigma^{-1} \left[\Sigma(\chi) \sin \frac{\alpha}{2} \right]. \end{aligned} \quad (29)$$

The fractional error made when using equation (28), $\frac{\Delta L}{L} = \frac{L_A - L}{L}$, is of course the same as in (25) for $\Delta z = 0$ since the extra factor $(1+z)$ cancels out. However, comparing equations (28) and (29) it is particularly easy to see that approximations such as (25) actually contain two approximations: (a) small angle and (b) neglect of curvature (for $k = \pm 1$). E.g. $\frac{\Delta L}{L}$ is independent of z in the flat case but is larger and varies with z for the open case.

ACKNOWLEDGMENTS

We thank C. Lineweaver for useful discussions.

REFERENCES

- Alcock C., Paczyński B., 1979, *Nat*, 281, 358
- Dinshaw N., Impey C. D., 1996, *ApJ*, 458, 73
- Fernández-Soto A., Barcons X., Carballo R., Webb J. K., 1995, *MNRAS*, 277, 235
- Halliday D., Resnick R., Walker J., 1993, *Fundamentals of Physics*. John Wiley & Sons, New York
- Kayser R., Helbig P., Schramm T., 1997, *A&A*, 318, 680
- Misner C. W., Thorne K. S., Wheeler J. A., 1973, *Gravitation*. W. H. Freeman and Company, San Francisco
- Osmer P. S., 1981, *ApJ*, 247, 762
- Peacock J. A., 1999, *Cosmological Physics*. Cambridge University Press, Cambridge
- Phillipps S., 1994, *MNRAS*, 269, 1077
- Popowski P. A., Weinberg D. H., Ryden B. S., Osmer P. S., 1998, *ApJ*, 498, 11
- Weinberg S., 1972, *Gravitation and Cosmolgy*. John Wiley & Sons, New York
- Williger G. M., Hazard C., Baldwin J. A., McMahon R. G., 1996, *ApJS*, 104, 145
- Yoshii Y., Peterson B. A., Takahara F., 1993, *ApJ*, 414, 431